

Interaction Report**Report ID:****Date Produced:**

28 May 2025

Antiretroviral Treatment**Co-medications**

Dolutegravir (DTG)

Cyanocobalamin (Vitamin B12) [in multivitamins]

This report lists the summaries of potential interactions (i.e. "red", "amber" and "yellow" classifications) for the drugs in the table above.

Interactions with a "green" or "grey" classification (i.e. no clinically significant interaction or no clear data) have been checked and are listed at the end of this report, but summaries are not shown.

For full details of all interactions, see www.hiv-druginteractions.org.

Description of the interactions

Potential clinically significant interaction - likely to require additional monitoring, alteration of drug dosage or timing of administration (AMBER)

Dolutegravir (DTG) + Cyanocobalamin (Vitamin B12) [in multivitamins]

Coadministration has not been studied. Cyanocobalamin undergoes enterohepatic recycling and is excreted almost entirely in the urine. No interaction is expected with vitamin B12 when given as a single agent (i.e. not in a multivitamin preparation) or as an injection. However, vitamin B12 can be found in multivitamin preparations; simultaneous coadministration of a multivitamin preparation decreased dolutegravir exposure by ~33%. Dolutegravir should be administered 2 hours before or 6 hours after medications containing polyvalent cations, such as multivitamin preparations. Medicinal products that reduce dolutegravir exposure (e.g. multivitamins) should be avoided in the presence of integrase class resistance.